

*Proceedings of XII International Conference on Selected Problems of Modern Physics,
Section I, Dubna, June 8–11, 2003.*

On the faster-than-light motions in electrodynamics

G. A. Kotel'nikov

RRC Kurchatov Institute, Moscow 123182, Russia

Abstract — A variant of electrodynamics is constructed in which the faster-than-light motions are possible.

The existence of faster-than-light motions (with velocities $v > 3 \cdot 10^{10}$ cm/sec) is the subject of discussion in modern physics.

It is in 1946–1948 when Blokhintsev already [1] paid attention to the possibility of formulating a field theory that permits the propagation of faster-than-light (superluminal) interactions outside the light cone. Kirzhnits (1954) [2] showed that a particle possessing the tensor of mass $M^i_k = \text{diag}(m_0, m_1, m_1, m_1)$ may move with the faster-than-light velocity if $m_0 > m_1$. Terletsky (1960) [3] introduced into theoretical physics the particles with imaginary masses moving faster-than-light. Feinberg (1967) [4] named these particles tachyons and described their main properties. Research on the superluminal tachyon motions opened up additional opportunities which were studied by many authors (hundreds publications), for example by Bilaniuk and Sudarshan (Sb. [4]), Kirzhnits and Sazonov (Sb. [4]), Recami [5], Mignani (Mon. [5]), Corben (Mon. [5]).

The publications are also known in which the violation of invariance of the speed of light is considered [6] - [12]. One can note, for example, Pauli monograph [6] with the elements of Ritz and Abraham theories; Logunov lectures on Relativity Theory [7]; Glashow paper [8] on the violation of Lorentz-invariance in astrophysics; publications [9] - [12] considering the violation of invariance of the speed of light in SR.

Below a version of the theory permitting faster-than-light motions of electromagnetic fields and charged particles with real masses is proposed as the continuation of such investigations.

Let us introduce space-time R^4 with the metric

$$ds^2 = (c_0^2 + v^2)dt^2 - dx^2 - dy^2 - dz^2 = (c_0'^2 + v'^2)dt'^2 - dx'^2 - dy'^2 - dz'^2 - invariant, \quad (1)$$

where t is the time, x, y, z are the spatial variables, v is the velocity of a particle being investigated, c_0 is the proper value of the speed of light. Due to homogeneity and isotropy of space, the velocity v does not depend on space-time variables. Let the proper value of the speed of light be invariant:

$$c_0 = c'_0 = 3 \cdot 10^{10} \text{ cm/sec.} \quad (2)$$

As a result, the common time may be introduced on the trajectory of movement of frame K' and the velocity of light c , corresponding to the velocity v : $dt = dt'_0 \rightarrow c = c_0 \sqrt{1 + v^2/c_0^2}$ ¹. Let us introduce also the new variables

$$x^0 = \int_0^t c d\tau = c_0 \int_0^t \sqrt{1 + v^2/c_0^2} d\tau, \quad x^\alpha = (x, y, z), \quad \alpha = 1, 2, 3, \quad (3)$$

and turn to Minkowski space M^4 with the metric

$$ds^2 = (dx^0)^2 - (dx^1)^2 - (dx^2)^2 - (dx^3)^2 - \text{invariant.} \quad (4)$$

The infinitesimal transformations, retaining the invariance of the form ds^2 are $dx'^i = L^i_k dx^k$, $i, k = 0, 1, 2, 3$, where L^i_k is the matrix of Lorentz group [14]. The corresponding integral transformations are

$$x'^0 = \frac{x^0 - \beta x^1}{\sqrt{1 - \beta^2}}; \quad x'^1 = \frac{x^1 - \beta x^0}{\sqrt{1 - \beta^2}}; \quad x'^{2,3} = x^{2,3}; \quad c' = c \frac{1 - \beta u^1}{\sqrt{1 - \beta^2}}. \quad (5)$$

They belong to the group of direct product $L_6 X S_1$, where L_6 is the Lorentz group, S_1 is the scale transformations group $c' = \gamma c$. One can say that these transformations act in the 5-space $V^5 = M^4 X V^1$, where V^1 is a subspace of the velocities of light c . The relationship between the partial derivatives of variables (t, x, y, z) and (x^0, x^1, x^2, x^3) are as follows:

$$\begin{aligned} \frac{\partial}{\partial t} &= \frac{\partial x^0}{\partial t} \frac{\partial}{\partial x^0} + \sum_\alpha \frac{\partial x^\alpha}{\partial t} \frac{\partial}{\partial x^\alpha} = c \frac{\partial}{\partial x^0}; \\ \frac{\partial}{\partial x} &= \frac{\partial x^0}{\partial x} \frac{\partial}{\partial x^0} + \sum_\alpha \frac{\partial x^\alpha}{\partial x} \frac{\partial}{\partial x^\alpha} = \left(\int_0^t \frac{\partial c}{\partial x} d\tau \right) \frac{\partial}{\partial x^0} + \frac{\partial}{\partial x^1}. \end{aligned} \quad (6)$$

The expressions for $\partial/\partial y$ and $\partial/\partial z$ are analogous to the expression $\partial/\partial x$. Below we restrict our study by the case, when the velocity of light in the range of interaction does not depend on the space variables. Then

$$\nabla c(x^0) = 0 \rightarrow \nabla c(t) = 0. \quad (7)$$

¹In the form of $c' = c(1 - \beta^2)^{1/2}$ this formula was obtained by Abraham before [6].

Some features of this study are:

1. As in SR, parameter $\beta = V/c$ in the present work is in the range $0 \leq \beta < 1$.
2. As in SR, the value dx^0 is the exact differential.
3. The time $x^0 = \int c dt$ is the functional of $c(t)$ in general case.
4. The property $\beta = \text{const}$ is compatible with $V(t)$, $c(t)$.

Keeping this in mind, let us construct a theory in M^4 and reflect it on the space R^4 . Following [13], we start from the expression for integral of action:

$$S = S_m + S_{mf} + S_f = -mc_0 \int ds - \frac{e}{c_0} \int A_i dx^i - \frac{1}{16\pi c_0} \int F_{ik} F^{ik} d^4x. \quad (8)$$

Here S_m is the action for a free particle; S_f is the action for a free electromagnetic field; S_{mf} is the action for interaction between a charge e and electromagnetic field; m is the mass of a particle, $j^i = (\rho, \rho \mathbf{u}^\alpha)$ [6] is the 4-vector of current density; $\mathbf{u}^\alpha = \mathbf{v}^\alpha/c$ is the 3-velocity of a particle. The meaning of the other values is standard. Accordingly to the construction, the action is Lorentz invariant and does not depend on the velocity of light c . As a result the action is also invariant with respect to the group of direct product L_6XS_1 . Lagrangian takes the form:

$$L = -mc_0 \sqrt{1 - u^2} + \frac{e}{c_0} (\mathbf{A} \cdot \mathbf{u} - \phi). \quad (9)$$

The generalized momentum $\mathbf{P}$ and generalized energy $\mathcal{H}$ are:

$$\begin{aligned} \mathbf{P} &= \frac{\partial L}{\partial \mathbf{u}} = \frac{mc_0 \mathbf{u}}{\sqrt{1 - u^2}} + \frac{e}{c_0} \mathbf{A} = \mathbf{p} + \frac{e}{c_0} \mathbf{A} = m\mathbf{v} + \frac{e}{c_0} \mathbf{A}, \\ \mathcal{H} &= \mathbf{P} \cdot \mathbf{u} - L = (mc_0 c + e\phi)/c_0. \end{aligned} \quad (10)$$

Here $\mathbf{p} = m\mathbf{v}$ is the momentum, $mc_0 c = mc_0^2 (1 + v^2/c_0^2)^{1/2} = \mathcal{E}$ is the energy, $T = mc_0^2 (c/c_0 - 1)$ is the kinetic energy of a particle. For a free particle they are integrals of motion, $\mathcal{E}$ and $\mathbf{p}$ may be united into the 4-momentum p^i (as in SR)

$$p^i = mc_0 u^i = \left(\frac{mc_0 c}{c_0}, mc u^\alpha \right) = \left(\frac{\mathcal{E}}{c_0}, m\mathbf{v} \right). \quad (11)$$

The components of the 4-momentum are related by the expressions:

$$p_i p^i = \frac{\mathcal{E}^2}{c_0^2} - \mathbf{p}^2 = m^2 c_0^2; \quad \mathbf{p} = \frac{\mathcal{E}}{c_0 c} \mathbf{v}; \quad \mathbf{p} = \frac{\mathcal{E}}{c_0 c} \mathbf{c}, \text{ if } m = 0, \mathbf{v} = \mathbf{c}. \quad (12)$$

One can see from here that the momentum of a particle with the zero mass $m = 0$ does not depend from the particle velocity $v = c$ and is only determined by the particle energy in accordance with $\mathbf{p} = \mathbf{n}\mathcal{E}/c_0$, $\mathbf{n} = \mathbf{c}/c$.

Next we start from the mechanical [13] and field [13,14] Lagrange equations:

$$\frac{d}{dx^0} \frac{\partial L}{\partial \mathbf{u}} - \frac{\partial L}{\partial \mathbf{x}} = 0; \quad \frac{\partial}{\partial x^k} \frac{\partial \mathcal{L}}{\partial (\partial A_i / \partial x^k)} - \frac{\partial \mathcal{L}}{\partial A_i} = 0. \quad (13)$$

Here L is the Lagrangian; $\mathcal{L} = -(1/c_0)A_i j^i - (1/16\pi c_0)F_{ik}F^{ik}$ is the density of the Lagrange function; $\partial(F_{ik}F^{ik})/\partial(\partial A_i/\partial x^k) = -4F^{ik}$ [13]. As a result, we find the equations of motion for a charged particle and electromagnetic field:

$$\begin{aligned} \frac{d\mathbf{p}}{dt} &= m_0 \frac{d\mathbf{v}}{dt} = \frac{c}{c_0} e \mathbf{E} + \frac{e}{c_0} \mathbf{v} \times \mathbf{H}; \\ \frac{d\mathcal{E}}{dt} &= e \mathbf{E} \cdot \mathbf{v} \rightarrow m_0 \frac{dc}{dt} = \frac{e}{c_0} \mathbf{v} \cdot \mathbf{E}. \end{aligned} \quad (14)$$

$$\begin{aligned} \nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{H}}{\partial t} &= 0; \quad \nabla \cdot \mathbf{E} = 4\pi\rho; \\ \nabla \times \mathbf{H} - \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} &= 4\pi\rho \frac{\mathbf{v}}{c}; \quad \nabla \cdot \mathbf{H} = 0, \end{aligned} \quad (15)$$

where $c(t) = c_0(1 + v^2/c_0^2)^{1/2} = c(0)[1 + (e/mc_0c(0)) \int_0^t \mathbf{v} \cdot \mathbf{E} d\tau]$, $\nabla c = 0$.

These equations, considered as the whole, form the set of nonlinear equations of electrodynamics. They permit the existence of faster-than-light motion of a particle with the real mass m , rest energy $\mathcal{E}_0 = mc_0^2$ and velocity

$$v = \sqrt{\mathcal{E}^2 - m^2 c_0^4} / mc_0 > c_0, \quad (16)$$

if the particle energy is $\mathcal{E} > \sqrt{2}\mathcal{E}_0$ [12]. For example, for electron $\sqrt{2}\mathcal{E}_0$ is 723 keV, the velocity of 1 GeV electron is $\sim 2000 c_0$. The $\mathcal{E}$ and $\mathbf{p}$ variations with time determine a new dynamics which goes to Newton one if $v^2/c_0^2 \ll 1$.

Explained in the framework of the constructed theory may be the Michelson experiment, Fizeau experiment, aberration of light, dilatation of life time of atmospheric μ - mesons, Doppler effect, known tests to check independence of the speed of light from the velocity of the light source, Compton effect, decay of unstable particles, creation of new particles in nuclear reactions, possible faster-than-light motions of nuclear reaction products. For example, in the case of Compton effect [15] we find from the law of energy-momentum conservation: $\hbar\omega = \hbar\omega' + mc_0^2 \left[\left(1 + v^2/c_0^2\right)^{1/2} - 1 \right]$; $(\hbar\omega/c_0) = (\hbar\omega'/c_0)\cos\theta + mv\cos\alpha$, $0 = (\hbar\omega'/c_0)\sin\theta - mv\sin\alpha$. Here $\hbar\omega$, $\hbar\omega'$ are the energies of incident and scattered γ quanta, mc_0^2 is the rest energy of electron, α and θ are the angles of scattering the electron and γ quantum respectively. As a result the angular distribution

of scattered γ quanta is $\omega' = \omega/[1 + \hbar\omega(1 - \cos\theta)/mc_0^2]$. (As in SR [15]). But the velocity of forward-scattered electron may exceed the speed of light c_0 : $v_e(\alpha = 0)^2 = \hbar\omega/mc_0 - 0.5c_0 > c_0$ if $\hbar\omega > 1.5mc_0^2$, which is in accordance with formula (16) and which differs from SR. The velocity of scattered γ quantum does not depend on the angle θ and is determined by mechanism of scattering (immediately, or in the act of absorption-emission by scattered electron).

As a result the theory has been constructed which is invariant with respect to the group of direct product L_6XS_1 . In accordance with [1] we may assume that the proposed theory may be useful in the field of quantum physics of extended particles, where the property of elementary nature should not contradict to the existence of the internal structure of a particle.

References

- [1] D. I. Blokhintsev, JETP, **16**, 480-482 (1946); **18**, 566-574 (1948).
- [2] D. A. Kirzhnits, JETP, **27**, 6-18 (1954).
- [3] Ya. P. Terletsky, DAN (Russ.), **133**, 329-332 (1960).
- [4] G. Feinberg, *Einstein Sbornik 1973* (M., Nauka, 1974), pp. 134-177.
- [5] E. Recami, *Relativity Theory and its Generalization. Astrophysics, Quanta and Relativity Theory* (M., Izdatelstvo Mir, 1982) pp. 53-128.
- [6] W. Pauli, *Theory of Relativity* (M.-L., Gostexizdat, 1947) pp. 29, 116.
- [7] A. A. Logunov, *Lectures on Fundamentals of Relativity Theory* (Moscow State University, 1982) pp. 25-50.
- [8] S. L. Glashow, Nucl. Phys. B (Proc. Suppl.), **70**, 180-184 (1999).
- [9] P. M. Rapier, IRE, **50**, 229-230 (1962).
- [10] J. P. Hsu and L. Hsu, Phys. Lett. A, **196**, 1-6 (1994).
- [11] A. Chubykalo and R. Smirnov-Rueda, Phys. Rev. E, **53**, 5373-5381 (1996).
- [12] G. A. Kotelnikov, J. of Russian Laser Research, **22**, 455-474 (2001).
- [13] L. D. Landau and E. M. Lifshitz. *The Theory of Field* (M., Fizmatgiz, 1973) pp. 68-71, 93-95, 110.
- [14] N. N. Bogoliubov and D. V. Shirkov. *Introduction to the Theory of Quantized Fields* (M., Nauka, 1973) pp. 16-17.
- [15] D. I. Blokhintsev, *Fundamentals of Quantum Mechanics* (M.-L., Gostexizdat, 1949) pp. 17-18.

$$^2v_e(\alpha = 0) = c_0(\hbar\omega/mc_0^2)[1 - mc_0^2/(2\hbar\omega + mc_0^2)] \approx \hbar\omega/mc_0 - 0.5c_0$$